



ITALPOLLINA
HELLO NATURE!

MYR

BORON




LEAF LIQUID INTEGRATORS

MYR RANGE is based on micronutrients and meso elements complexed with vegetal peptides with the aim of prevent and/or cure nutritional deficiencies and improve yield and fruit quality.

MYR RANGE guarantees the highest efficiency and a quick absorption, without losses due to insolubilization. It is the result of a careful research on the mechanisms of vegetal absorption in order to prevent phenomena of phytotoxicity caused by micronutrients.

The peptides of **MYR** are quickly absorbed and then incorporated in the lymph : they can carry out their functions in stressed plants in a few hours, allowing a remarkable saving of energy and permitting the recovery of the optimal conditions for development.

Oligonutritional complexes **MYR** have a stable structure also in alkaline environment, which prevents the formation of salts or insoluble hydroxides.



MYR RANGE is powered by **LISIVEG**, an exclusive production system designed by the R&D of Italpollina, based on the **enzymatic hydrolysis of vegetal proteins**. It allows to obtain **PEPTIDES** that, unlike individual free amino acids, have a more prominent, assured and **natural biostimulant effect**. **LISIVEG®** lies its strength in the innovative treatment technology that **preserves even the weakest components** (as Tryptophan) and maintains its activity unaltered.



MYR BORON

MYR BORON increases the pollen fertility, formation and quality of fruits, it improves photosynthesis and fruit setting, reducing fruit drop. Boron is easily assimilable and mobile inside the plants.

Organic Nitrogen (N)	4 %
Boron (B) water soluble	5 %
Organic carbon (C)	18 %
pH (1:5)	7,5
Specific weight	1,24 kg/L

THE ADVANTAGES

- Increase the pollen fertility
- Improvement of photosynthesis and fruit setting, reducing fruit drop
- Increases the formation and quality of fruits
- Antistress and nutritional action



USE & DOSE

All dosages provided are in standard conditions, dosages and modes of application are to be determined on the basis of the requirements and the nutritional status of the crop. We recommend the application in the coolest hours of the day (morning or evening) to obtain maximum absorption by the leaves and the best effectiveness.

MYR range can be mixed with the most common phytosanitary products and fertilizers on the market. Before using we recommend testing on a small surface for compatibility with other products.

Liquid fertilizer for foliar application (on leaves).

Crops	Dose recommended		Stage
	Dosage per ha	Dosage per hl	
Vegetables	1,5 L	150 mL	Every 15 days
Orchards	2 L	200 mL	2-3 applications before flowering period
Soft fruits	2 L	200 mL	Every 15 days starting in preflowering period
Vineyards, pineapple, bananas, kiwi	2 L	200 mL	2-3 applications before flowering period
Sunflower	3 L	1,5 L*	Limit tractor crossing 7-8 leaves
Sugar beet	3 L	1,5 L*	2 applications to cover crops needs

* based on 200 L/ha of spray solution

MYR

BORON



PACKAGING
1 - 5 - 20 L



*Suitable for organic farming
according to EC REG. 834/2007*

ITALPOLLINA S.p.A.
Località Casalmenini, 10
37010 Rivoli Veronese (VR) Italia
Tel +39.045.6205511
info@italpollina.com
www.italpollina.com

Follow us   