






## **LEAF LIQUID INTEGRATORS**

**MYR** RANGE is based on micronutrients and meso elements complexed with vegetal peptides with the aim of prevent and/or cure nutritional deficiencies and improve yield and fruit quality.

**MYR** RANGE guarantees the highest efficiency and a quick absorption, without losses due to insolubilization. It is the result of a careful research on the mechanisms of vegetal absorption in order to prevent phenomena of phytotoxicity caused by micronutrients.

The peptides of MYR are quickly absorbed and then incorporated in the lymph: they can carry out their functions in stressed plants in a few hours, allowing a remarkable saving of energy and permitting the recovery of the optimal conditions for development.

Oligonutritional complexes **MYR** have a stable structure also in alkaline environment, which prevents the formation of salts or insoluble hydroxides.



MYR RANGE is powered by LISIVEG, an exclusive production system designed by the R&D of



Italpollina, based on the **enzymatic hydrolysis of vegetal**proteins. It allows to obtain **PEPTIDES** that, unlike individual free amino acids, have a more prominent, assured and **natural** 

**biostimulant effect. LISIVEG®** lies its strength in the innovative treatment technology that **preserves even the weakest components** (as Tryptophan) and maintains its activity unaltered.





## **MYR MAGNESIUM**

MYR MAGNESIUM improves photosynthesis and sugars in the fruits, it also prevents dessication of the grape spine and apple leaf drop. It improves fruit firmness and quality and color of the leaves. It has an anti-stress and nutritional effect.

Organic Nitrogen (N)	3 %
Magnesium oxyde (MgO) water soluble	5 %
Organic carbon (C)	16 %
pH (1:5)	5
Specific weight	1,27 kg/L

## THE ADVANTAGES

- Increase the tolerance of plants to environmental stresses
- Improvement of photosynthesis and fruit setting, reducing fruit drop
- Ouick nutritional action
- Chlorosis Limitation



## **USE & DOSE**

All dosages provided are in standard conditions, dosages and modes of application are to be determined on the basis of the requirements and the nutritional status of the crop. We recommend the application in the coolest hours of the day (morning or evening) to obtain maximum absorption by the leaves and the best eff ectiveness.

MYR range can be mixed with the most common phytosanitary products and fertilizers on the market. Before using we recommend testing on a small surface for compatibility with other products.

Liquid fertilizer for foliar application (on leaves).

Crops	Dose recommended		Stage	
	Dosage per ha	Dosage per hl		
Vegetables	3 L	300 mL	Around flowering stage to cover Mg needs of plants - 3 applications	
Orchards, Soft fruits	3 L	300 mL	Post flowering stage - 3 applications	
Vineyards	5 L	300 mL	From fruit setting to veraison - 3 applications	
Ananas	5 L	300 mL	In case of need	
Bananas	5 L	300 mL	Before flowering stage to cover important needs of plants - 3 applications	
kiwi	5 L	300 mL	Post flowering stage - 3 applications	
Nursery		300 mL	Every week	



**PACKA SING** 1 – 5 – 20 <u>L</u>



ITALPOLLINA S.p.A. Località Casalmenini, 10 37010 Rivoli Veronese (VR) Italia Tel +39.045.6205511 info@italpollina.com www.italpollina.com

Follow us f Tube in



