

10-6-4






NUTRITIONAL BALANCE FOR QUALITY HARVESTS

NUTRIMYR NPK 10-6-4 thanks to its high content of nitrogen (organic and ureic) and the balanced presence of phosphorus, potassium and micronutrients, improves plant growth and helps to overcome abiotic stress (excessive heat, cold, drought etc..)

ENHANCED FOLIARS FOR A COMPLETE AND BALANCED NUTRITION



COMPOSITION

Total nitrogen (N)	10 %
> Organic nitrogen (N)	2 %
> Ureic nitrogen (N)	8 %
Phosphorus pentoxide (P₂O₅) total	6 %
Potassium oxide (K ₂ O) water soluble	4 %
Boron (B) water soluble	
Manganese (Mn) chelated with EDTA water soluble	
Molybdenum (Mo) water soluble	0,02 %
Zinc (Zn) chelated with EDTA water soluble	0,05 %
Organic carbon (C)	12 %
pH (1:5)	4
Specific weight	1,18 kg/L

THE ADVANTAGES

- Improve of quality, consistency and shelf life of fruits
- Increase of yield and quality of the harvest
- Stimulation of the formation of chlorophyll
- Improvement of the photosynthetic activity
- Antistress action

NUTRIMYR RANGE is powered by **LISIVEG®**, an exclusive production system designed by the R&D



of Italpollina, based on the enzymatic hydrolysis of vegetal proteins. It allows to obtain PEPTIDES that, unlike individual free amino acids, have a more prominent, assured and natural biostimulant effect. LISIVEG®

lies its strength in the innovative treatment technology that preserves even the weakest components (as Tryptophan) and maintains its activity unaltered.



USE AND DOSE

All dosages provided are in standard conditions, dosages and modes of application are to be determined on the basis of the requirements and the nutritional status of the crop.

We recommend the application in the coolest hours of the day (morning or evening) to obtain maximum absorption by the leaves and the best eff ectiveness.

NUTRIMYR range can be mixed with the most common phytosanitary products and fertilizers on the market. Before using we recommend testing on a small surface for compatibility with other products.

Liquid fertilizer for foliar application (on leaves).

Crops	Dose recommended	Stage
	Dosage per ha	
Vineyards	3-5 L	Repeating 2-3 times (until the formation of clusters)
Orchards	3-5 L	Every 15 days after petals fall - 4 to 6 applications
Horticulture	3-5 L	Every 15 days from the beginning of the production cycle - 3 to 5 applications



Tel +39.045.6205511 info@italpollina.com www.italpollina.com



